

Neutrophil-to-lymphocyte ratio in patients with gram-negative sepsis admitted to intensive care unit

Naser Gharebaghi, Mohammad Amin Valizade Hasanloei, Alireza Mehdizadeh Khalifani, Shiva Pakzad, Durna Lahooti

Urmia University of Medical Sciences, Urmia, Iran

Abstract

Background: Several studies have investigated the role of the neutrophil-to-lymphocyte ratio (N/L ratio) in the early diagnosis of infection with controversial results. We assessed this ratio in the early diagnosis of gram-negative sepsis cases admitted to intensive care unit (ICU).

Methods: Patients with gram-negative sepsis admitted to general intensive care unit were enrolled. The following data was recorded: demographic characteristics; APACHE II score; duration of mechanical ventilation and ICU length of stay; average neutrophil and lymphocyte count, along with their ratio, on the first, second and third day of hospitalisation; and the mortality rate.

Results: One hundred and thirty nine patients were studied during a period of one year with a mean age 68.29 ± 17.4 years. One hundred and thirty six patients received mechanical ventilation, with a mean duration of 13.85 ± 2.07 days. The mean white blood cell (WBC) count on the first day of hospitalisation was 11776.04 ± 5789.60 , while on the second and third days this was 12446 ± 2101.16 and 13241.01 ± 7077.17 , respectively. The N/L ratio on the day of hospitalisation was 11.38 ± 1.02 , and 11.98 ± 0.99 and 12.94 ± 1.3 on the second and third days, respectively. Moreover, 46.8 percent of patients died during the study.

From applying Student's t-test, significant differences were observed between two groups of patients (discharged and deceased) in terms of their APACHE II score, neutrophil count, WBC and N/L ratio on the second and third days of hospitalisation ($P < 0.05$).

Conclusion: Blood cell analysis and N/L ratio can be used as a predictor for the severity of gram-negative sepsis along with other diagnostic procedures.

Anestezjologia Intensywna Terapia 2019, tom 51, nr 1, 11–16

Key words: sepsis, gram-negative; neutrophil; lymphocyte; neutrophil-to-lymphocyte ratio; biomarkers

Sepsis is one of the oldest syndromes in clinical medicine dating back to the time of Hippocrates [1] and it is one of the major causes of morbidity and mortality among patients in intensive care units [2], with more than 750,000 sepsis cases each year being hospitalised in the United States [3]. Earlier definitions of sepsis and septic shock defined them as cardiovascular dysfunction following an infection that has no other cause [5] and suggested that sepsis is the body's response to the damaged tissues and organs caused by the infection [6].

The new definition of sepsis, based on the Third International Consensus Definitions of Sepsis and Septic Shock in 2016, define it as a life-threatening state and organ dysfunction caused by a dysregulated host response to infection. In this regard, the immune modulatory system has a vital role and affects the cardiovascular, nervous, autonomic, humeral, metabolic and coagulation systems. For quicker diagnosis, instead of using SIRS and SOFA scores, qSOFA scores have been proposed, which included mental impairment, a res-

piratory rate greater than or equal to 22 breaths per minute, and systolic pressure less than or equal to 100 mmHg. These criteria are considered as predictive of the mortality in patients with sepsis: with one of the above criteria, the mortality rate would be less than 3%, and with all three criteria present, this risk would be more than 20% [7–9].

Although sepsis and septic shock have mostly bacterial causes, fungi and parasites may also cause it, while in one third of cases no infectious cause is detected. Symptoms include as follows: fever (hypothermia in 10–15% of cases); leukocytosis (occasionally leucopenia); lactic acidosis (blood lactate higher than 2 mmol L⁻¹); continued arterial pressure drop and the need for a vasopressor with sufficient fluid replacement; low urine output; decreased peripheral blood flow; and altered mental status. The most common characteristic of septic shock is its hyperkinetic pattern, which is associated with a high cardiac output [10].

Currently, accessible, cheap, and fast biomarkers are used to predict inflammatory and infectious reactions. Numerous studies have evaluated the role of inflammatory tests, including the neutrophil-to-lymphocyte ratio (N/L ratio), in the pre-determination of infections and cancers in various parts of the body and which have yielded contradictory results. During the acute phase of inflammation, particularly as a result of bacterial infection, environmental exposure [11] and some cancers, neutrophils are one of the first inflammatory responders of cells to migrate towards the site of inflammation [12, 13]. Therefore, the N/L ratio is used as a marker of subclinical inflammation [14] and tumours [15–18].

One study regarding the diagnosis of adult complicated acute appendicitis, performed in 2012, did not show sufficient accuracy for this test [19]. Regarding another study on pulmonary infection caused by *Escherichia coli*, although showing an ability to predict disease severity, its low test ratio did not allow for rejecting possibilities of bacteraemia or acute sepsis [20]. Our study evaluated the predictive power of the neutrophil-to-lymphocyte ratio as a fast marker in patients with gram-negative sepsis.

METHODS

This cross sectional study was approved by ethics committee of Urmia University of Medical Sciences and conducted in the GICUs (General Intensive Care Units) of Imam and Ayatollah Taleghani hospitals (tertiary educational hospitals) in Urmia during 2015. Due to the prevalence of approximately 15–25% of sepsis patients in the ICU, nearly 200 patients with sepsis are hospitalised in these two GICUs each year. All non-surgical patients admitted to a GICU who developed gram-negative sepsis during their hospitalisation, or those who were hospitalised with gram-negative

sepsis, were included in the study. Patients with Glasgow Coma Scale – 3 and brain death were excluded, along with all surgical, trauma and cancer patients. Diagnosis was based on clinical and vital signs, having a positive blood culture, or other accompanying tests, such as a white blood cell count and differential, the erythrocyte sedimentation rate, C-reactive protein, procalcitonin levels, as well as a diagnosis by an infectious disease specialist according to the new definition of sepsis-3 [7]. The neutrophil and lymphocyte counts, and their ratio, were recorded on the first, second and third days. Furthermore, the APACHE score II, demographic information, duration of hospitalisation, mortality rate in the ICU, duration of mechanical ventilation, as well as the types of antibiotics used were recorded from the medical information ward.

A statistical analysis was performed using SPSS (Ver. 20, 2010). Student's t-test was used to compare quantitative variables with a normal distribution while the Chi-square test was used for qualitative variables with a normal distribution. $P < 0.05$ was considered statistically significant.

RESULTS

In this cross-sectional study, 139 patients with gram-negative sepsis and a mean age of 68.29 ± 17.4 years were enrolled. Among them, 72 (51.8%) were female and 67 (48.2%) were male. The mean APACHE II score was 17.38 ± 3.27 . *Kelbsiella sp.* (33.81%), *Escherichia coli* (27.33%), *Pseudomonas sp.* (15.1%), *Enterobacter sp.* (12.23%), *Acinetobacter sp.* (10.07%), *Serratia sp.* (1.435%) were commonly found in blood cultures of patients. Eight patients (5.8%) received only one antibiotic, 76 (54.7%) received two, 43 (30.9%) three, 11 (7.9%) four, while one patient (0.7%) received five antibiotics.

The mean length of stay in the ICU was 16.79 ± 1.83 days. Only three patients did not receive supportive mechanical ventilation and the mean duration of mechanical ventilation was 13.58 ± 2.07 days for the remaining patients. Tables 1 and 2 show the demographic characteristics for discharged and deceased patients.

Although seventy four patients (53.2%) with an APACHE II score of 16.23 ± 2.17 were discharged after a full recovery, unfortunately 65 patients (46.8%) with an APACHE II score of 21.08 ± 1.82 died during the study Student's t-test ($P = 0.02$).

The mean white blood cell count (WBCs) during hospitalisation was 11776.04 ± 5789.60 . However, mean WBCs on the second and third days were 12446 ± 2101.16 and 13241.01 ± 7077.17 , respectively. Table 3 shows a comparison between discharged and deceased patients in terms of WBCs during the study.

Table 4 shows a comparison between discharged and deceased patients in terms of neutrophil percentage during the study.

Table 1. Demographic and baseline parameters

Baseline parameters		Discharged	Deceased	P-value
Age (years)		65.81 ± 13.2	68.7 ± 18.34	0.06
Gender	Male	42	30	0.78
	Female	38	29	
ICU stay (days)		16.4 ± 1.23	22.17 ± 0.78	0.03
Body mass index (kg m ⁻²)		27.91 ± 4.6	26.13 ± 3.58	0.07
Fasting blood sugar (mg dL ⁻²)		95.28 ± 4.3	101.05 ± 3.18	0.04
Hb ₁ C		5.1 ± 0.23	6.1 ± 0.33	0.04
Mortality rate		57.56%	42.44%	0.03

Table 2. Characteristics of disease in discharged and deceased patients

Disease	Total	Discharged	Deceased
Cerebrovascular accident	36	20	16
Chronic obstructive pulmonary disease	21	14	7
Aspiration pneumonia and pneumonia	13	10	3
Lung cancer	5	2	3
Venous thromboembolism	14	8	6
End-stage renal disease	31	16	15
Rheumatologic disorder	13	7	6
Oncologic disorders	6	3	3
Total	139	80	59

Table 3. Comparison between discharged and deceased patients in term of white blood cells during the study

	Outcome	Neutrophil	P-value
First day	Discharged	11226.08 ± 5149.39	0.23
	Deceased	12402.15 ± 8425.31	
Second day	Discharged	11170.68 ± 4731.60	0.001
	Deceased	13898.92 ± 5152.54	
Third day	Discharged	11043.24 ± 4157.58	0.0001
	Deceased	15743.08 ± 8737.24	

Table 4. Comparison between discharged and deceased patients in terms of neutrophil percentage during the study

	Outcome	Neutrophil	P-value
First day	Discharged	81.54 ± 9.24	0.2
	Deceased	83.63 ± 9.92	
Second day	Discharged	81.84 ± 8.25	0.008
	Deceased	85.69 ± 8.55	
Third day	Discharged	80.64 ± 9.50	0.01
	Deceased	85.05 ± 10.31	

Table 5 shows a comparison between discharged and deceased patients in terms of lymphocyte percentage during the study.

The N/L ratio on the first day of hospitalisation was 11.38 ± 1.02, while on the second and third days this was assessed as 11.98 ± 0.99 and 12.94 ± 1.3, respectively. Table 6 shows

Table 5. Comparison between discharged and deceased patients in terms of lymphocyte percentage during the study

	Outcome	Neutrophil	P-value
First day	Discharged	14.21 ± 8.65	0.17
	Deceased	12.17 ± 9.12	
Second day	Discharged	13.23 ± 7.40	0.04
	Deceased	10.62 ± 8.35	
Third day	Discharged	14.29 ± 8.19	0.002
	Deceased	10.03 ± 7.96	

Table 6. Comparison of discharged and deceased patients in terms of neutrophil-to-lymphocyte ratio (N/L ratio)

	Outcome	N/L ratio	P-value
First day	Discharged	9.88 ± 10.54	0.12
	Deceased	13.09 ± 1.68	
Second day	Discharged	9.38 ± 7.78	0.004
	Deceased	14.95 ± 1.80	
Third day	Discharged	9.17 ± 1.37	0.002
	Deceased	17.23 ± 2.20	

Table 7. Comparison of discharged and deceased patients in terms of blood culture.

Blood culture	Discharged	Deceased	P-value
<i>Klebsiella</i>	26	21	0.1
<i>Escherichia coli</i>	25	13	0.05
<i>Pseudomonas</i>	11	10	0.1
<i>Enterobacter</i>	10	7	0.6
<i>Acinetobacter</i>	7	7	1
<i>Serratia</i>	1	1	1

a comparison of discharged and deceased patients in terms of the neutrophil-to-lymphocyte ratio.

Table 7 shows a comparison of discharged and deceased patients in terms of blood culture.

DISCUSSION

The results of our study showed that there were significant differences between the two groups of patients concerned (discharged and deceased patients) in term of APACHEII scores, WBCs and neutrophil counts and N/L ratios.

Many studies have attempted to introduce accessible, cheap and fast biomarkers, including the WBC count or N/L ratio, for the quick detection of sepsis. Zahorec reported that the N/L ratio may express the severity of disease and can be regularly used in the ICU at 6, 12 and 24-hour intervals [21]. Similarly, our patients who died due to gram-negative sepsis had had a higher N/L ratio in serial tests (second and third day) compared with patients who had fully recovered.

Because we had excluded patients with trauma or surgery and malignancy, we considered the reason of the elevation of the neutrophil count and N/L ratio was sepsis.

Several studies have investigated the N/L ratio in infectious patients, with contradictory results [19, 20, 22, 23].

Hosseinpour *et al.* [19] investigated the diagnostic value of this ratio in acute adult appendicitis, and claimed that despite a significant difference between the positive and negative pathology patients, this test lacks sufficient accuracy in diagnosing acute and symptomatic appendicitis. One year later, Ljungstorm *et al.* [20] claimed that the N/L ratio could be a marker of disease severity for *Escherichia coli* infection, and the higher this ratio, the higher the possibility of severe sepsis. On the other hand, a low ratio never precludes the possibility of bacteraemia or severe sepsis. Similarly, our study showed that patients with a poor prognosis (the deceased group with a high APACHE II score) had had a higher N/L ratio and that this ratio, along with other tests, could be used to predict disease severity. However,

we claim that this ratio cannot be used by itself to diagnose gram-negative sepsis.

The importance of sepsis and its early diagnosis and treatment and implications in all areas are emphasised [24, 25].

In a retrospective study, De Jager *et al.* [26] compared patients with positive and negative blood cultures. They indicated that the N/L ratio and lymphocytopenia bacteraemia are better predictors than the routine parameters in an emergency care setting. Similar to their study, we showed that an increased neutrophil-to-lymphocyte ratio during the first 3 days of admission indicates sepsis severity and an increased rate of mortality. An increase in this ratio is either due to an increase in the number of neutrophils or a decrease in the number of lymphocytes or both, and it may be concluded that lymphopenia may have existed in our study. We have neglected to examine them since they did not fall within the remit of our study.

Liu *et al.* [27] indicated that an increased N/L ratio was independently associated with an unfavourable clinical prognosis in patients with sepsis. Median N/L ratio levels were significantly higher in patients who died than in survivors. Our study confirms these findings.

It has been also claimed that the N/L ratio can be used as a novel non-invasive marker to predict the occurrence of hospital-acquired bacterial infections in decompensated cirrhotic patients [28].

Akilli *et al.* [29] suggested that the N/L ratio in critically ill patients is an important indicator of short- and long-term mortalities: thus, hospital mortality and 6-month mortality were associated with a higher N/L ratio. This was a prospective cohort study, and although patients were about ten years older than patients in our study, they were similar in terms of gender.

Hwang *et al.* [30] stated that the initial N/L ratio measured on admission to an emergency department was independently associated with 28-day mortality in patients with severe sepsis and septic shock, and that the changes in this ratio may provide an important prognostic marker.

Gurol *et al.* [31] examined 1,468 patients with bacteraemia and sepsis. They concluded that the total white cell and neutrophil counts have historically been used as markers of infection. They also believed that an N/L ratio ≥ 5 indicated a more convenient marker than CRP because of its ability to detect bacterial infections at a lower cost. As with these studies, our study, though with a different sample size, showed that a high neutrophil/lymphocyte ratio on the second and third days correlated with an increase in the mortality of hospitalised patients with gram-negative sepsis in the ICU. It also showed that this ratio may be used, along other with diagnostic procedures and markers of disease severity, for predicting the outcome of these patients. Thus, in our

deceased patients the N/L ratio was more than 10 and in discharged patients less than 10.

This study suffered from some inevitable limitations. Incomplete patient records in the hospitals that we encountered was one of the shortcomings of the study that decreased the number of the cases studied. Furthermore, selecting patients with only gram-negative sepsis and excluding cases with gram-positive was another limitation. Finally, the duration of the study and the sample size of the population was limited in our study. It is recommended that physicians conduct a study with both gram-positive and gram-negative sepsis over a longer period of time and with a large sample size in the near future.

CONCLUSION

According to the results of our study, an analysis of the serial CBC and neutrophil/lymphocyte ratio, along other with tests, can provide a fast method for establishing disease identification, severity and outcome in sepsis.

ACKNOWLEDGEMENTS

1. Source of funding: none.
2. Conflict of interest: none.

References:

1. Angus DC, Vander Poll T. Severe sepsis and septic shock. *N Engl J Med.* 2013; 349: 840–851.
2. Kaukonen KM, Bailey M, Pilcher D, et al. Systemic inflammatory response syndrome criteria in defining severe sepsis. *N Engl J Med.* 2015; 372(17): 1629–1638, doi: [10.1056/NEJMoa1415236](https://doi.org/10.1056/NEJMoa1415236), indexed in Pubmed: [25776936](https://pubmed.ncbi.nlm.nih.gov/25776936/).
3. Kellum JA, Pike F, Yealy DM, et al. and the Protocol-based Care for Early Septic Shock Investigators (ProCESS) Investigators, ProCESS Investigators. A randomized trial of protocol-based care for early septic shock. *N Engl J Med.* 2014; 370(18): 1683–1693, doi: [10.1056/NEJMoa1401602](https://doi.org/10.1056/NEJMoa1401602), indexed in Pubmed: [24635773](https://pubmed.ncbi.nlm.nih.gov/24635773/).
4. Seymour C, Liu V, Iwashyna T, et al. Assessment of Clinical Criteria for Sepsis. *JAMA.* 2016; 315(8): 762–774, doi: [10.1001/jama.2016.0288](https://doi.org/10.1001/jama.2016.0288).
5. Shankar-Hari M, Phillips GS, Levy ML, et al. Sepsis Definitions Task Force. Developing a New Definition and Assessing New Clinical Criteria for Septic Shock: For the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA.* 2016; 315(8): 775–787, doi: [10.1001/jama.2016.0289](https://doi.org/10.1001/jama.2016.0289), indexed in Pubmed: [26903336](https://pubmed.ncbi.nlm.nih.gov/26903336/).
6. Rhodes A, Evans LE, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Crit Care Med.* 2017; 45(3): 486–552, doi: [10.1097/CCM.0000000000002255](https://doi.org/10.1097/CCM.0000000000002255), indexed in Pubmed: [28098591](https://pubmed.ncbi.nlm.nih.gov/28098591/).
7. Singer M, Deutschman CS, Seymour CW, et al. Sepsis Definitions Task Force. Assessment of Clinical Criteria for Sepsis: For the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA.* 2016; 315(8): 762–774, doi: [10.1001/jama.2016.0288](https://doi.org/10.1001/jama.2016.0288), indexed in Pubmed: [26903335](https://pubmed.ncbi.nlm.nih.gov/26903335/).
8. Van der Steen M, Verhage R, Singer M, et al. Overview of the Third International Consensus Definitions for Sepsis and Shock (sepsis – 3). *Netherlands Journal of Critical Care.* 2016; 6–9.
9. Fleischmann C, Scherag A, Adhikari NKJ, et al. International Forum of Acute Care Trialists. Assessment of Global Incidence and Mortality of Hospital-treated Sepsis. Current Estimates and Limitations. *Am J Respir Crit Care Med.* 2016; 193(3): 259–272, doi: [10.1164/rccm.201504-0781OC](https://doi.org/10.1164/rccm.201504-0781OC), indexed in Pubmed: [26414292](https://pubmed.ncbi.nlm.nih.gov/26414292/).
10. Vincent JL. Septic shock. In: Vincent JL, Abraham E, Moore FA, Kochanek PM, Fink MP, eds. *Textbook of critical care 7th ed.*. Elsevier, Oxford 2017: 843–848.

11. Jacobs L, Nawrot TS, de Geus B, et al. Subclinical responses in healthy cyclists briefly exposed to traffic-related air pollution: an intervention study. *Environ Health*. 2010; 9: 64, doi: [10.1186/1476-069X-9-64](https://doi.org/10.1186/1476-069X-9-64), indexed in Pubmed: [20973949](https://pubmed.ncbi.nlm.nih.gov/20973949/).
12. Waugh D, Wilson C. The interleukin-8 pathway in cancer. *Clinical Cancer Research*. 2008; 14(21):6735–6741, doi: [10.1158/1078-0432.ccr-07-4843](https://doi.org/10.1158/1078-0432.ccr-07-4843).
13. De Larco JE, Wuertz BRK, Furcht LT. The potential role of neutrophils in promoting the metastatic phenotype of tumors releasing interleukin-8. *Clin Cancer Res*. 2004; 10(15): 4895–4900, doi: [10.1158/1078-0432.CCR-03-0760](https://doi.org/10.1158/1078-0432.CCR-03-0760), indexed in Pubmed: [15297389](https://pubmed.ncbi.nlm.nih.gov/15297389/).
14. Wang X, Zhang G, Jiang X, et al. Neutrophil to lymphocyte ratio in relation to risk of all-cause mortality and cardiovascular events among patients undergoing angiography or cardiac revascularization: a meta-analysis of observational studies. *Atherosclerosis*. 2014; 234(1): 206–213, doi: [10.1016/j.atherosclerosis.2014.03.003](https://doi.org/10.1016/j.atherosclerosis.2014.03.003), indexed in Pubmed: [24681815](https://pubmed.ncbi.nlm.nih.gov/24681815/).
15. Wang J, Jia Y, Wang N, et al. The clinical significance of tumor-infiltrating neutrophils and neutrophil-to-CD8+ lymphocyte ratio in patients with resectable esophageal squamous cell carcinoma. *J Transl Med*. 2014; 12: 7, doi: [10.1186/1479-5876-12-7](https://doi.org/10.1186/1479-5876-12-7), indexed in Pubmed: [24397835](https://pubmed.ncbi.nlm.nih.gov/24397835/).
16. Wei Y, Jiang YZ, Qian WH. Prognostic role of NLR in urinary cancers: a meta-analysis. *PLoS One*. 2014; 9(3): e92079, doi: [10.1371/journal.pone.0092079](https://doi.org/10.1371/journal.pone.0092079), indexed in Pubmed: [24642859](https://pubmed.ncbi.nlm.nih.gov/24642859/).
17. Young CA, Murray LJ, Karakaya E, et al. The Prognostic Role of the Neutrophil-to-Lymphocyte Ratio in Oropharyngeal Carcinoma Treated with Chemoradiotherapy. *Clin Med Insights Oncol*. 2014; 8: 81–86, doi: [10.4137/CMO.S15476](https://doi.org/10.4137/CMO.S15476), indexed in Pubmed: [25057245](https://pubmed.ncbi.nlm.nih.gov/25057245/).
18. Xiao WK, Chen D, Li SQ, et al. Prognostic significance of neutrophil-lymphocyte ratio in hepatocellular carcinoma: a meta-analysis. *BMC Cancer*. 2014; 14: 117, doi: [10.1186/1471-2407-14-117](https://doi.org/10.1186/1471-2407-14-117), indexed in Pubmed: [24559042](https://pubmed.ncbi.nlm.nih.gov/24559042/).
19. Hosseinpour M, Arbabi M, Nazadi S, et al. Evaluation of diagnostic value of neutrophil-lymphocyte ratio in acute and complicated appendicitis of adults utility of neutrophil lymphocyte ratio in acute complicated cholecystitis. *Journal of Surgery*. 2012(3): 39.
20. Ljungstrom LR, Jacobsson G, Andersson R. Neutrophil-lymphocyte count ratio as a biomarker of severe sepsis in *Escherichia coli* infections in adults. *Critical Care*. 2013; 17(Suppl 2): P25, doi: [10.1186/cc11963](https://doi.org/10.1186/cc11963).
21. Zahorec R. Ratio of neutrophil to lymphocyte counts—rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy*. 2001; 102(1): 5–14, indexed in Pubmed: [11723675](https://pubmed.ncbi.nlm.nih.gov/11723675/).
22. Saliccioli JD, Marshall DC, Pimentel MAF, et al. The association between the neutrophil-to-lymphocyte ratio and mortality in critical illness: an observational cohort study. *Crit Care*. 2015; 19: 13, doi: [10.1186/s13054-014-0731-6](https://doi.org/10.1186/s13054-014-0731-6), indexed in Pubmed: [25598149](https://pubmed.ncbi.nlm.nih.gov/25598149/).
23. Sarran Y, Mengi T, Yilmaz B, et al. Can N-L Ratio predict diseases severity and mortality in sepsis and septic shock patients? *Acta medica mediterranea*. 2018: 877–882.
24. Yoldas H, Karagoz I, Ogun MN, et al. Novel Mortality Markers for Critically Ill Patients. *J Intensive Care Med*. 2018 [Epub ahead of print]: 885066617753389, doi: [10.1177/0885066617753389](https://doi.org/10.1177/0885066617753389), indexed in Pubmed: [29334832](https://pubmed.ncbi.nlm.nih.gov/29334832/).
25. Arit KS, Rukka ABS, Wahyuni S. Comparison of NLR and procalcitonin parameters in sepsis patients treated in intensive care unit. *Journal of Medical Sciences*. ; 2017: 17–21.
26. de Jager CPC, van Wijk PTL, Mathoera RB, et al. Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia better than conventional infection markers in an emergency care unit. *Crit Care*. 2010; 14(5): R192, doi: [10.1186/cc9309](https://doi.org/10.1186/cc9309), indexed in Pubmed: [21034463](https://pubmed.ncbi.nlm.nih.gov/21034463/).
27. Liu X, Shen Y, Wang H, et al. Prognostic significance of neutrophil-to-lymphocyte ratio in patients with sepsis: a prospective observational study. *Mediators Inflamm*. 2016; 2016: 8191254, doi: [10.1155/2016/8191254](https://doi.org/10.1155/2016/8191254), indexed in Pubmed: [27110067](https://pubmed.ncbi.nlm.nih.gov/27110067/).
28. Cai YJ, Dong JJ, Dong JZ, et al. Neutrophil-lymphocyte ratio predicts hospital-acquired bacterial infections in decompensated cirrhosis. *Clin Chim Acta*. 2017; 469: 201–207, doi: [10.1016/j.cca.2017.04.011](https://doi.org/10.1016/j.cca.2017.04.011), indexed in Pubmed: [28412195](https://pubmed.ncbi.nlm.nih.gov/28412195/).
29. Akilli NB, Yortanlı M, Mutlu H, et al. Prognostic importance of neutrophil-lymphocyte ratio in critically ill patients: short- and long-term outcomes. *Am J Emerg Med*. 2014; 32(12): 1476–1480, doi: [10.1016/j.ajem.2014.09.001](https://doi.org/10.1016/j.ajem.2014.09.001), indexed in Pubmed: [25264245](https://pubmed.ncbi.nlm.nih.gov/25264245/).
30. Hwang SY, Shin TG, Jo IkJ, et al. Neutrophil-to-lymphocyte ratio as a prognostic marker in critically-ill septic patients. *Am J Emerg Med*. 2017; 35(2): 234–239, doi: [10.1016/j.ajem.2016.10.055](https://doi.org/10.1016/j.ajem.2016.10.055), indexed in Pubmed: [27806894](https://pubmed.ncbi.nlm.nih.gov/27806894/).
31. Gürol G, Çiftçi İH, Terizi HA, et al. Are there standardized cutoff values for neutrophil-lymphocyte ratios in bacteremia or sepsis? *J Microbiol Biotechnol*. 2015; 25(4): 521–525, indexed in Pubmed: [25341467](https://pubmed.ncbi.nlm.nih.gov/25341467/).

Adres do korespondencji:

Mohammad Amin Valizade
Urmia Medical Faculty, Patient Safety Research Centre
Urmia, Iran
e-mail: aminvalizade@yahoo.com

Otrzymano: 5.05.2018 r.

Zaakceptowano: 6.03.2019 r.